arXiv:1112.4706v1 [math.DS] 20 Dec 2011

# ON THE NUMBER OF FIXED POINTS OF SOFIC FLIP SYSTEMS

YOUNG-ONE KIM AND SIEYE RYU

Abstract. In the case when $X$ is a sofic shift and $\varphi : X \to X$ is a homeomorphism such that $\varphi^2 = \mathrm{id}_X$ and $\varphi\sigma_X = \sigma_X^{-1}\varphi$, the number of points in $X$ that are fixed by $\sigma_X^m$ and $\sigma_X^n\varphi$, $m = 1, 2, \ldots$, $n \in \mathbb{Z}$, is expressed in terms of a finite number of square matrices: The matrices are obtained from Krieger's joint state chain of a sofic shift which is conjugate to $X$.

## 1. Introduction

Let $(X, T)$ be an invertible topological dynamical system. A homeomorphism $F : X \to X$ is called a *flip map* (or simply a *flip*) for $(X, T)$ if

$$FT = T^{-1}F \quad \text{and} \quad F^2 = \mathrm{id}_X. \tag{1.1}$$

In this case, we call the triple $(X, T, F)$ a flip system. Two flip systems $(X, T, F)$ and $(X', T', F')$ are said to be *conjugate* if there is a homeomorphism $\theta : X \to X'$ such that

$$\theta \circ T = T' \circ \theta \quad \text{and} \quad \theta \circ F = F' \circ \theta.$$

We call the homeomorphism $\theta$ a *conjugacy* from $(X, T, F)$ to $(X', T', F')$.

If $m$ is a positive integer, the number of periodic points in $(X, T)$ of period $m$ is denoted by $p_m(T)$:

$$p_m(T) = |\{x \in X : T^m(x) = x\}|.$$

Similarly, if $F$ is a flip for $(X, T)$, $m$ is a positive integer and $n$ is an integer, then $p_{m,n}(T, F)$ will denote the number of points in $X$ that are fixed by $T^m$ and $T^nF$:

$$p_{m,n}(T, F) = |\{x \in X : T^m(x) = T^nF(x) = x\}|.$$

It is obvious that if the flip systems $(X, T, F)$ and $(X', T', F')$ are conjugate, then $p_m(T) = p_m(T')$ and $p_{m,n}(T, F) = p_{m,n}(T', F')$ for all $m$ and $n$. From (1.1), it follows that

$$p_{m,n}(T, F) = p_{m,n+m}(T, F) = p_{m,n+2}(T, F).$$

Hence $p_{m,n}(T, F) = p_{m,0}(T, F)$ if $m$ and $n$ are even; $p_{m,n}(T, F) = p_{m,1}(T, F)$ if $m$ is even and $n$ is odd; and $p_{m,n}(T, F) = p_{m,0}(T, F)$ for all $n$ if $m$ is odd.

It is well known that if $X$ is a shift of finite type, then there is a square matrix $A$ whose entries are non-negative integers such that

$$p_m(\sigma_X) = \mathrm{tr}(A^m) \qquad (m = 1, 2, \cdots). \tag{1.2}$$

It is also well known that if $X$ is a sofic shift, then there are square matrices $A_1, A_2, \cdots, A_r$ whose entries are integers such that

$$p_m(\sigma_X) = \sum_{k=1}^{r} (-1)^{k+1} \mathrm{tr}(A_k^m) \qquad (m = 1, 2, \cdots).$$

Suppose that the sequence $\{(p_m(T))^{1/m}\}$ is bounded. Then the Artin-Mazur zeta function $\zeta_T$ of $(X, T)$ is defined by

$$\zeta_T(t) = \exp\left(\sum_{m=1}^{\infty} \frac{p_m(T)}{m} t^m\right);$$

and the results mentioned above imply that if $X$ is a shift of finite type, then $\zeta_{\sigma_X}$ is the reciprocal of a polynimial, and if $X$ is a sofic shift, then $\zeta_{\sigma_X}$ a rational function.

In [KLP], Y.-O. Kim, J. Lee and K.K.Park introduced a zeta function $\zeta_{T,F}$ of a flip system which is given by

$$\zeta_{T,F}(t) = \zeta_T(t^2)^{1/2} \exp(G_{T,F}(t)),$$

where

$$G_{T,F}(t) = \sum_{m=1}^{\infty} \left(p_{2m-1,0}(T, F) t^{2m-1} + \frac{p_{2m,0}(T, F) + p_{2m,1}(T, F)}{2} t^{2m}\right).$$

The function $G_{T,F}$ will be called the *generating function* of $(X, T, F)$.

In the case when $X$ is a shift of finite type and $\varphi$ is a flip for $(X, \sigma_X)$, a formula for $p_{m,n}(\sigma_X, \varphi)$ which is similar to (1.2) is known. In order to present it, we need some notation: If M is a matrix $\mathcal{S}[M]$ will denote the sum of the entries of M, that is,

$$\mathcal{S}[M] = \sum_{I,J} M_{IJ}$$

and $M^\Delta$ will denote the matrix whose diagonal entries are identical with those of $M$ but the other entries are equal to 0, that is,

$$(M^\Delta)_{IJ} = \begin{cases} M_{IJ} & \text{if } I = J, \\ 0 & \text{otherwise.} \end{cases}$$

For instance, $\mathrm{tr}(M) = \mathcal{S}[M^\Delta]$, whenever $M$ is a square matrix.

**Theorem A.** *If $X$ is a shift of finite type and $\varphi$ is a flip for $(X, \sigma_X)$, then there are zero-one square matrices $A$ and $J$ such that*

$$\begin{aligned} p_m(\sigma_X) &= \mathrm{tr}(A^m), \\ p_{2m,0}(\sigma_X, \varphi) &= \mathcal{S}[J^\Delta A^m J^\Delta], \\ p_{2m,1}(\sigma_X, \varphi) &= \mathcal{S}[(JA)^\Delta A^{m-1} (AJ)^\Delta] \quad \textit{and} \\ p_{2m-1,0}(\sigma_X, \varphi) &= \mathcal{S}[J^\Delta A^{m-1} (AJ)^\Delta] \qquad (m = 1, 2, \cdots). \end{aligned}$$

**Remark.** As it is proved in [KLP], Theorem A implies that if $X$ is a shift of finite type and $\varphi$ is a flip for $(X, \sigma_X)$, then the generating function of $(X, \sigma_X, \varphi)$ is a rational function.

In this paper, we extend Theorem A to the case when $X$ is a sofic shift.

**Theorem B.** *If $X$ is a sofic shift and $\varphi$ is a flip for $(X, \sigma_X)$, then there are square matrices $A_k, B_k$ and $J_k$, $k = 1, 2, \cdots, r$, whose entries are integers such that*

$$\begin{aligned}
p_m(\sigma_X) &= \sum_{k=1}^{r} (-1)^{k+1} \operatorname{tr}(A^m), \\
p_{2m,0}(\sigma_X, \varphi) &= \sum_{k=1}^{r} (-1)^{k+1} \mathcal{S}\big[J_k^{\Delta} B_k^m J_k^{\Delta}\big], \\
p_{2m,1}(\sigma_X, \varphi) &= \sum_{k=1}^{r} (-1)^{k+1} \mathcal{S}\big[(J_k A_k)^{\Delta} B_k^{m-1} (A_k J_k)^{\Delta}\big] \quad \textit{and} \\
p_{2m-1,0}(\sigma_X, \varphi) &= \sum_{k=1}^{r} (-1)^{k+1} \mathcal{S}\big[J_k^{\Delta} B_k^{m-1} (A_k J_k)^{\Delta}\big] \qquad (m = 1, 2, \cdots).
\end{aligned}$$

The same method as in the proof of Theorem 3.2 in [KLP] gives the following.

**Corollary.** *If $X$ is a sofic shift and $\varphi$ is a flip for $(X, \sigma_X)$, then the generating function of $(X, \sigma_X, \varphi)$ is a rational function.*

We prove Theorem B in Section 2. In our proof of the theorem, Krieger's joint state chain plays a crucial role and it is described in Section 3. We conclude the paper with an example and some remarks(Section 4).

## 2. Proof of Theorem B

We start with some preliminaries. Suppose $\mathcal{A}$ is a finite alphabet and $\tau : \mathcal{A} \to \mathcal{A}$ satisfies $\tau^2 = \mathrm{id}_{\mathcal{A}}$. Then the map $\varphi_\tau : \mathcal{A}^{\mathbb{Z}} \to \mathcal{A}^{\mathbb{Z}}$ defined by

$$\varphi_\tau(x)_i = \tau(x_{-i})$$

is a flip for $(\mathcal{A}^{\mathbb{Z}}, \sigma)$. If $X$ is a shift space over $\mathcal{A}$ and $\varphi_\tau(X) = X$, the restriction of $\varphi_\tau$ to $X$ is a flip for $(X, \sigma_X)$. Conversely, if $X$ is a shift space over $\mathcal{A}$, $\varphi$ is a flip for $(X, \sigma_X)$ and $\varphi$ satisfies

$$x, x' \in X \text{ and } x_0 = x'_0 \quad \Rightarrow \quad \varphi(x)_0 = \varphi(x')_0,$$

then there is a map $\tau : \mathcal{A} \to \mathcal{A}$ such that $\tau^2 = \mathrm{id}_{\mathcal{A}}$, $\varphi_\tau(X) = X$ and $(\varphi_\tau)|_X = \varphi$. In this case, we say that $\varphi$ is a *one-block flip* and we call the restriction of $\tau$ to $\mathcal{B}_1(X)$ the *symbol map* of $\varphi$.

If $A$ is a zero-one $\mathcal{A} \times \mathcal{A}$ matrix, $\mathsf{X}_A$ will denote the topological Markov chain determined by $A$:

$$\mathsf{X}_A = \{x \in \mathcal{A}^{\mathbb{Z}} : \forall i \in \mathbb{Z} \;\; A(x_i, x_{i+1}) = 1\}.$$

We denote by $\sigma_A$ the restriction of the shift map to $\mathsf{X}_A$. Let $A$ be such a matrix. Then we have $\varphi_\tau(\mathsf{X}_A) = \mathsf{X}_A$ whenever

$$A(a, b) = A(\tau(b), \tau(a)) \qquad (a, b \in \mathcal{A}) \tag{2.1}$$

holds. If we define the $\mathcal{A} \times \mathcal{A}$ matrix $J$ by

$$J(a, b) = \begin{cases} 1 & \text{if } \tau(a) = b, \\ 0 & \text{otherwise,} \end{cases} \tag{2.2}$$

then $J$ is a symmetric permutation matrix, $J^2 = I$ and the condition (2.1) is equivalent to the equation

$$JA = A^{\mathsf{T}} J. \tag{2.3}$$

Conversely, if $J$ is a zero-one $\mathcal{A} \times \mathcal{A}$ matrix such that $J^2 = I$, then it is a symmetric permutation matrix and hence there is a unique map $\tau : \mathcal{A} \to \mathcal{A}$ such that (2.2) holds and consequently $\tau^2 = \mathrm{id}_{\mathcal{A}}$. In this case, the flip $\varphi_\tau$ will also be denoted by $\varphi_J$. If (2.3) holds, the flip $\varphi_J$ to $\mathsf{X}_A$ is a flip for $(\mathsf{X}_A, \sigma_A)$. We denote the restriction by $\varphi_{J,A}$.

The following proposition will be proved in the next section.

**Proposition 2.1.** *Suppose $X$ is a sofic shift and $\varphi$ is a flip for $(X, \sigma_X)$. Then there are a finite set $\mathcal{A}$, a map $\mathcal{L} : \mathcal{A} \to \mathcal{B}_1(X)$ and zero-one $\mathcal{A} \times \mathcal{A}$ matrices $A$ and $J$ having the following properties:*
(1) *$\mathcal{L}_\infty : \mathsf{X}_A \to X$ is a factoring.*
(2) *$\mathcal{L}_\infty$ has no graph diamonds.*
(3) *$JA = A^{\mathsf{T}} J$ and $J^2 = I$.*
(4) *$\mathcal{L}_\infty \circ \varphi_{J,A} = \varphi \circ \mathcal{L}_\infty$.*
(5) *If $\delta \in \{0, 1\}$, $x \in X$ and $\sigma_X^\delta \varphi(x) = x$, then there is a $y \in \mathsf{X}_A$ such that $\mathcal{L}_\infty(y) = x$ and $\sigma_A^\delta \varphi_{J,A}(y) = y$.*

In the rest of this section, we suppose that $X$ is a sofic shift, $\varphi$ is a flip for $(X, \sigma_X)$ and the system $(\mathcal{A}, \mathcal{L}, A, J)$ is as in the above proposition. We denote by $\tau$ the map of $\mathcal{A}$ onto itself such that $J(a, \tau(a)) = 1$ for all $a \in \mathcal{A}$.

If $\pi$ is a permutation of a finite set, its sign will be denoted by $\mathrm{sgn}(\pi)$. When $<$ is a linear order of $\mathcal{A}, S \subset \mathcal{A}$ and $f : S \to \mathcal{A}$ is one-to-one, we set

$$\mathrm{sgn}_<(f) = (-1)^{N(<, f)},$$

where

$$N(<, f) = |\{(a, b) \in S \times S : a < b \text{ and } f(b) < f(a)\}|.$$

If $g : f(S) \to \mathcal{A}$ is one-to-one again, we have

$$\mathrm{sgn}_<(g \circ f) = \mathrm{sgn}_<(g)\, \mathrm{sgn}_<(f).$$

We also have $\mathrm{sgn}_<(f) = \mathrm{sgn}(f)$ wherever $f$ is a permutation of a subset of $\mathcal{A}$.

From here on, we fix a linear order $<$ of $\mathcal{A}$ and write

$$\mathrm{sgn}(f) = \mathrm{sgn}_<(f)$$

for arbitrary one-to-one function $f$ from a subset of $\mathcal{A}$ into $\mathcal{A}$.

Let $k$ be a positive integer $\leq |\mathcal{A}|$. We set

$$\mathcal{A}_k = \{S \subset \mathcal{A} : |S| = k \text{ and } |\mathcal{L}(S)| = 1\};$$

and for $S_1, S_2 \in \mathcal{A}_k$ we denote by $F(S_1, S_2)$ the set of one-to-one functions $f : S_1 \to S_2$ such that $A(a, f(a)) = 1$ for all $a \in S_1$. Define the $\mathcal{A}_k \times \mathcal{A}_k$

matrices $A_k, B_k$ and $J_k$ by

$$A_k(S_1,S_2) = \sum_{f\in F(S_1,S_2)} \operatorname{sgn}(f),$$

$$B_k(S_1,S_2) = \begin{cases} 1 & \text{if } F(S_1,S_2)\neq\varnothing, \\ 0 & \text{otherwise,}\end{cases}$$

and

$$J_k(S_1,S_2) = \begin{cases} \operatorname{sgn}(\tau|_{S_1}) & \text{if } \tau(S_1)=S_2, \\ 0 & \text{otherwise.}\end{cases}$$

It is well known that

$$p_m(\sigma_X) = \sum_{k=1}^{|\mathcal{A}|} (-1)^{k+1}\operatorname{tr}(A_k^m) \qquad (m=1,2,\cdots). \tag{2.4}$$

Hence the first formula in Theorem B holds. A proof of (2.4) is found in Section 6.4 of [LM](See also [B] and [M].) and our proof of the other formulas in Theorem B is a 'modification' of it. In order to explain the modification, we give an outline of the proof of (2.4).

Let $m$ be a positive integer. We put

$$P(m) = \{x\in X : \sigma_X(x)=x\},$$

so that $p_m(\sigma_X) = |P(m)|$. Suppose $x\in P(m)$. Since $\mathcal{L}_\infty : \mathsf{X}_A \to X$ is a factoring with no graph diamonds and since $\sigma_X^m(x)=x$, $\mathcal{L}_\infty^{-1}(x)$ is a nonempty finite subset of $\mathsf{X}_A$ and the resriction of $\sigma_A^m$ to $\mathcal{L}_\infty^{-1}(x)$ is a permutation of $\mathcal{L}_\infty^{-1}(x)$. Let $\mathcal{C}(m;\,x)$ denote the set of all subsets of $\mathcal{L}_\infty^{-1}(x)$ that are fixed by $\sigma_A^m$:

$$\mathcal{C}(m;\,x) = \{E\subset\mathcal{L}_\infty^{-1}(x) : \sigma_A^m(E)=E\}.$$

The following lemma is proved in section 6.4 of [LM].

**Lemma 2.2.** *Let $\pi$ be a permutation of a nonempty finite set $F$ and $\mathcal{C} = \{E\subset F : \pi(E)=E\}$. Then*

$$\sum_{E\in\mathcal{C}\setminus\{\varnothing\}} (-1)^{|E|+1}\operatorname{sgn}(\pi|_E) = 1.$$

With the help of this lemma, we obtain

$$p_m(\sigma_X) = \sum_{x\in P(m)}\ \sum_{E\in\mathcal{C}(m;\,x)\setminus\{\varnothing\}} (-1)^{|E|+1}\operatorname{sgn}(\sigma_A^m|_E). \tag{2.5}$$

For $k=1,2,\cdots,|\mathcal{A}|$ let $X_k$ denote the topological Markov chain determined by the zero-one $\mathcal{A}_k\times\mathcal{A}_k$ matrix $B_k$:

$$X_k = \{z\in\mathcal{A}_k^{\mathbb{Z}} : \forall i\in\mathbb{Z}\ F(z_i,z_{i+1})\neq\varnothing\}.$$

We denote the shift map of $X_k$ by $\sigma_k$ and put

$$P(m;\,k) = \{z\in X_k : \sigma_k^m(z)=z\}.$$

If $z\in P(m;\,k)$, then $|F(z_i,z_{i+1})|=1$ for all $i$, because $\mathcal{L}_\infty : \mathsf{X}_A\to X$ has no graph diamonds; hence $A_k(z_i,z_{i+1}) = 1$ or $-1$ for all $i$. In this case, we write

$$\operatorname{sgn}(z) = A_k(z_0,z_1)A_k(z_1,z_2)\cdots A_k(z_{m-1},z_m).$$

With this notation, we have

$$\sum_{z\in P(m;\,k)} \operatorname{sgn}(z) = \operatorname{tr}(A_k^m) \qquad (k = 1, 2, \cdots, |\mathcal{A}|). \tag{2.6}$$

Let $P_0$ denote the map $\mathsf{X}_A \ni y \mapsto y_0 \in \mathcal{A}$. Suppose $x \in P(m)$. Since $\mathcal{L}_\infty : \mathsf{X}_A \to X$ has no graph diamonds and $x$ is periodic, the restriction of $P_0$ to $\mathcal{L}_\infty^{-1}(x)$ is one-to-one. In particular, $|\mathcal{L}_\infty^{-1}(x)| \le |\mathcal{A}|$. If $E \in \mathcal{C}(m;\, x) \setminus \{\varnothing\}$ and $k = |E|$, then $1 \le k \le |\mathcal{A}|$, $P_0(\sigma_A^i(E)) \in \mathcal{A}_k$ for all $i$ and

$$\big(P_0(\sigma_A^i(E))\big)_{i\in\mathbb{Z}} \in P(m;\, k).$$

We denote the point $(P_0(\sigma_A^i(E)))_{i\in\mathbb{Z}}$ by $\Phi(E)$. With this notation, we have

$$\operatorname{sgn}(\sigma_A^m|_E) = \operatorname{sgn}(\Phi(E)), \qquad (E \in \mathcal{C}(m,\, x) \setminus \{\varnothing\}). \tag{2.7}$$

Finally, it is easy to check that the map

$$\Phi : \bigcup_{x\in P(m)} \mathcal{C}(m;\, x) \setminus \{\varnothing\} \to \bigcup_{k=1}^{|\mathcal{A}|} P(m;\, k)$$

is a one-to-one correspondence. Hence (2.4) follows from(2.5), (2.6)and(2.7).

In our proof of the other formulas in Theorem B, we need the following modification of Lemma 2.2.

**Lemma 2.3.** *Let $\pi$ be a permutation of a finite set $F$. Suppose $\mathcal{C} \subset 2^F$ and $G \subset F$ satisfy the following:*
*(1) $\pi(E) = E$ for every $E \in \mathcal{C}$.*
*(2) If $E_1, E_2 \in \mathcal{C}$, then $E_1 \cup E_2, E_1 \setminus E_2 \in \mathcal{C}$.*
*(3) $G \ne \varnothing$ and $\pi(G) = G$.*
*(4) If $E \subset G$ and $\pi(E) = E$, then $E \in \mathcal{C}$.*
*Then*

$$\sum_{E\in\mathcal{C}\setminus\{\varnothing\}} (-1)^{|E|+1} \operatorname{sgn}(\pi|_E) = 1. \tag{2.8}$$

*Proof.* The assumptions imply that $\mathcal{C}$ has at least two distinct elements, namely $G$ and $\varnothing$. For convenience, we set $\operatorname{sgn}(\pi|_\varnothing) = 1$. With this notation, (2.8) is equivalent to

$$\sum_{E\in\mathcal{C}} (-1)^{|E|} \operatorname{sgn}(\pi|_E) = 0.$$

If we put

$$\mathcal{N} = \{E \in \mathcal{C} : E \subset G\} \text{ and } \mathcal{R} = \{E \in \mathcal{C} : E \cap G = \varnothing\},$$

then (2) implies that

$$\mathcal{N} \times \mathcal{R} \ni (E_1, E_2) \mapsto E_1 \cup E_2 \in \mathcal{C}$$

is a one-to-one correspondence. If $(E_1, E_2) \in \mathcal{N} \times \mathcal{R}$, then $E_1 \cap E_2 = \varnothing$, and hence

$$(-1)^{|E_1\cup E_2|} \operatorname{sgn}(\pi|_{E_1\cup E_2}) = (-1)^{|E_1|} \operatorname{sgn}(\pi|_{E_1})\, (-1)^{|E_2|} \operatorname{sgn}(\pi|_{E_2}).$$

From (3) and Lemma 2.2, we have

$$\sum_{E\subset G,\, \pi(E)=E} (-1)^{|E|} \operatorname{sgn}(\pi|_E) = 0;$$

and from (1) and (4), we have

$$\{E \subset G : \pi(E) = E\} = \mathcal{N}.$$

Therefore

$$\begin{aligned}\sum_{E \in \mathcal{C}} (-1)^{|E|} \operatorname{sgn}(\pi|_E) &= \sum_{(E_1, E_2) \in \mathcal{N} \times \mathcal{R}} (-1)^{|E_1 \cup E_2|} \operatorname{sgn}(\pi|_{E_1 \cup E_2}) \\ &= \sum_{E_1 \in \mathcal{N}} (-1)^{|E_1|} \operatorname{sgn}(\pi|_{E_1}) \sum_{E_2 \in \mathcal{R}} (-1)^{|E_2|} \operatorname{sgn}(\pi|_{E_2}) \\ &= 0.\end{aligned}$$

□

**Remark.** The condition that $G \neq \varnothing$ is crucial. Otherwise we cannot apply Lemma 2.2.

Now, suppose that $N$ is a positive integer and $\delta \in \{0, 1\}$. We put

$$P(N, \delta) = \{x \in X : \sigma_X^N(x) = \sigma_X^\delta \, \varphi(x) = x\}$$

so that $p_{N,\delta}(\sigma_x, \varphi) = |P(N, \delta)|$; and for $x \in P(N, \delta)$ we put

$$\mathcal{C}(N, \delta;\, x) = \{E \subset \mathcal{L}_\infty^{-1}(x) : \sigma_A^N(E) = \sigma_A^\delta \, \varphi_{J,A}(E) = E\}.$$

We have $P(N, \delta) \subset P(N)$ and for $x \in P(N, \delta)$ we have $\mathcal{C}(N, \delta;\, x) \subset \mathcal{C}(N;\, x)$.

**Lemma 2.4.** *We have*

$$p_{N,\delta}(\sigma_X, \varphi) = \sum_{x \in P(N,\delta)} \sum_{E \in \mathcal{C}(N,\delta;\, x) \setminus \{\varnothing\}} (-1)^{|E|+1} \operatorname{sgn}(\sigma_k^N|_E).$$

*Proof.* Let $x \in P(N, \delta)$ and put

$$G(x) = \{y \in \mathcal{L}_\infty^{-1}(x) : \exists n \in \mathbb{Z} \;\; \sigma_A^\delta \, \varphi_{J,A}(y) = \sigma_A^{nN}(y)\}.$$

It follows from (5) in Proposition 2.1 that $G(x) \neq \varnothing$. It is then straightforward to check that the conditions in Lemma 2.3 are satisfied with $F = \mathcal{L}_\infty^{-1}(x)$, $\pi = \sigma_A^N|_{\mathcal{L}_\infty^{-1}(x)}$, $\mathcal{C} = \mathcal{C}(N, \delta;\, x)$ and $G = G(x)$. Hence we obtain

$$\sum_{E \in \mathcal{C}(N,\delta;\, x) \setminus \{\varnothing\}} (-1)^{|E|+1} \operatorname{sgn}(\sigma_A^N|_E) = 1 \qquad (x \in P(N, \delta)),$$

and the result follows. □

Suppose $1 \leq k \leq |\mathcal{A}|$ and let us consider the topological Markov chain $X_k$ again. If $S \in \mathcal{A}_k$, we have $\tau(S) \in \mathcal{A}_k$. We also have

$$B_k(S_1, S_2) = B_k(\tau(S_2), \tau(S_1)) \qquad (S_1, S_2 \in \mathcal{A}_k).$$

Hence the map $\varphi_k : X_k \to X_k$ defined by

$$\varphi_k(z)_i = \tau(z_{-i})$$

is a flip for $(X_k, \sigma_k)$. We put

$$P(N, \delta;\, k) = \{z \in X_k : \sigma_k^N(z) = \sigma_k^\delta \, \varphi_k(z) = z\}.$$

Again we have $P(N,\delta;\,k) \subset P(N;\,k)$ for every $k$. Moreover, if $x \in P(N,\delta)$, $E \in \mathcal{C}(N,\delta;\,x) \setminus \{\varnothing\}$ and $k = |E|$, then $1 \le k \le |\mathcal{A}|$ and $\Phi(E) \in P(N,\delta;\,k)$; and, as before, the map

$$\Phi : \bigcup_{x \in P(N,\delta)} \mathcal{C}(N,\delta;\,x) \setminus \{\varnothing\} \to \bigcup_{k=1}^{|\mathcal{A}|} P(N,\delta;\,k)$$

is a one-to-one correspondence.

If $E \in \mathcal{C}(N,\delta;\,x) \setminus \{\varnothing\}$ for some $x \in P(N,\delta)$, then (2.7) implies that

$$\operatorname{sgn}(\sigma_A^N|_E) = \operatorname{sgn}(\Phi(E)).$$

By Lemma 2.4, we have

$$p_{N,\delta}(\sigma_X,\varphi) = \sum_{k=1}^{|\mathcal{A}|} (-1)^{k+1} \sum_{z \in P(N,\delta;\,k)} \operatorname{sgn}(z).$$

Therefore the following lemma completes our proof of Theorem B.

**Lemma 2.5.** *Let* $1 \le k \le |\mathcal{A}|$. *Then we have*

$$\sum_{z \in P(N,\delta;\,k)} \operatorname{sgn}(z) = \begin{cases} \mathcal{S}\big[J_k^{\Delta} B_k^m J_k^{\Delta}\big] & \text{if } N = 2m \text{ and } \delta = 0, \\ \mathcal{S}\big[(J_k A_k)^{\Delta} B_k^{m-1} (A_k J_k)^{\Delta}\big] & \text{if } N = 2m \text{ and } \delta = 1, \\ \mathcal{S}\big[J_k^{\Delta} B_k^m (A_k J_k)^{\Delta}\big] & \text{if } N = 2m+1 \text{ and } \delta = 0. \end{cases}$$

*Proof.* We prove the last one only. The others are similarly proved. Let $m$ be a non-negative integer. We have to show that

$$\sum_{z \in P(2m+1,0;\,k)} \operatorname{sgn}(z) = \mathcal{S}\big[J_k^{\Delta} B_k^m (A_k J_k)^{\Delta}\big].$$

For $S \in \mathcal{A}_k$ we write $\tau(S) = S^*$. Thus $J_k(S_1,S_2) \neq 0$ if and only if $S_1^* = S_2$.

Let $S_0, S_1, \cdots, S_m \in \mathcal{A}_k^{m+1}$. We denote the product

$$J_k(S_0,S_0) B_k(S_0,S_1) \cdots B_k(S_{m-1},S_m) A_k(S_m,S_m^*) J_k(S_m^*,S_m)$$

by $\Pi(S_0 S_1 \cdots S_m)$. If $m = 0$,

$$\Pi(S_0) = J_k(S_0,S_0) A_k(S_0,S_0^*) J_k(S_0^*,S_0).$$

The expression $\mathbb{E}(S_0 S_1 \cdots S_m)$ will denote the unique point $z \in \mathcal{A}_k^{\mathbb{Z}}$ such that

$$z_0 z_1 \cdots z_{2m} = S_0 S_1 \cdots S_m S_m^* \cdots S_1^*$$

and

$$z_{i+2m+1} = z_i \qquad (i \in \mathbb{Z}).$$

If $m = 0$, then $\mathbb{E}(S_0)$ is the point in $\mathcal{A}_k^{\mathbb{Z}}$ such that $z_i = S_0$ for all $i$.

We have

$$\mathcal{S}\big[J_k^{\Delta} B_k^m (A_k J_k)^{\Delta}\big] = \sum_{S_0 S_1 \cdots S_m \in \mathcal{I}} \Pi(S_0 S_1 \cdots S_m),$$

where

$$\mathcal{I} = \{S_0 S_1 \cdots S_m \in \mathcal{A}_k^{m+1} : \Pi(S_0 S_1 \cdots S_m) \neq 0\},$$

and it is straightforward to check that the map

$$\mathcal{I} \ni S_0 S_1 \cdots S_m \mapsto \mathbb{E}(S_0 S_1 \cdots S_m) \in P(2m+1, 0;\,k)$$

is well-defined, one-to-one and onto. It remains only to show that

$$\Pi(S_0S_1\cdots S_m) = \operatorname{sgn}(\mathbb{E}(S_0S_1\cdots S_m))$$

holds for every $S_0S_1\cdots S_m \in \mathcal{I}$.

Let $S_0S_1\cdots S_m \in \mathcal{I}$. We have to show that

$$\begin{aligned}&\Pi(S_0S_1\cdots S_m)\\ &= A_k(S_0,S_1)\cdots A_k(S_{m-1},S_m)A_k(S_m,S_m^*)\cdots A_k(S_1^*,S_0).\end{aligned} \tag{2.9}$$

If we put $z = \mathbb{E}(S_0S_1\cdots S_m)$, then $|A_k(z_i,z_{i+1})| = 1$ for all $i$, and hence

$$\begin{aligned}&\Pi(S_0S_1\cdots S_m)\\ &= J_k(S_0,S_0)\big(A_k(S_0,S_1)\cdots A_k(S_{m-1},S_m)\big)^2 A_k(S_m,S_m^*)J_k(S_m^*,S_m).\end{aligned}$$

We also have $S_0^* = S_0$, because $J_k(S_0,S_0) \neq 0$.

It is easy to show that

$$J_k(S,\mathcal{T}) = J_k(\mathcal{T},S) \text{ and } J_k(S,S^*)A_k(S,\mathcal{T}) = A_k(\mathcal{T}^*,S^*)J_k(\mathcal{T},\mathcal{T}^*)$$

holds for all $S,\mathcal{T} \in \mathcal{A}_k$. Thus

$$\begin{aligned}&J_k(S_0,S_0)A_k(S_0,S_1)\cdots A_k(S_{m-1},S_m)J_k(S_m^*,S_m)\\ =\ &A_k(S_1^*,S_0^*)\cdots A_k(S_m^*,S_{m-1}^*)J_k(S_m,S_m^*)J_k(S_m^*,S_m)\\ =\ &A_k(S_m^*,S_{m-1}^*)\cdots A_k(S_1^*,S_0).\end{aligned}$$

Hence (2.9) follows. □

## 3. Krieger's Joint State Chain(Proof of Proposition 2.1)

We start with some notation. Suppose $\mathcal{A}$ is a finite alphabet, $j,k \in \mathbb{Z}$ and $j \le k$. If $x \in \mathcal{A}^{\mathbb{Z}}$, the expression $x_{[j,\,k]}$ will denote the restriction of $x$ to the set $[j,\,k] = \{i \in \mathbb{Z} : j \le i \le k\}$. Thus $x_{[j,\,k]} \in \mathcal{A}^{[j,\,k]}$. If $E \subset \mathcal{A}^{\mathbb{Z}}$, we also write $E_{[j,\,k]} = \{x_{[j,\,k]} : x \in E\}$. We will use similar notation for other subsets of $\mathbb{Z}$ as well.

The following lemma implies that every flip for a shift dynamical system is conjugate to a one-block one.

**Lemma 3.1.** *Suppose $X$ is a shift space and $\varphi$ is a flip for $(X,\,\sigma_X)$. Then there are a finite set $\mathcal{A}$, a map $\Phi : \mathcal{A} \to \mathcal{B}_1(X)$, a shift space $Y$ over $\mathcal{A}$ and a one-block flip $\psi$ for $(Y,\,\sigma_Y)$ such that $\Phi_\infty$ is a conjugacy from $(Y,\,\sigma_Y,\,\psi)$ to $(X,\,\sigma_x,\,\varphi)$.*

*Proof.* Since $\varphi$ is (uniformly) continuous, there is a non-negative integer $N$ such that

$$x,x' \in X \text{ and } x_{[-N,\,N]} = x'_{[-N,\,N]} \Rightarrow \varphi(x)_0 = \varphi(x')_0.$$

Put

$$\mathcal{A} = \{(a,b) : \exists x \in X \text{ s.t. } a = x_0 \text{ and } b = \varphi(x)_0\}$$

and define the maps $\Phi : \mathcal{A} \to \mathcal{B}_1(X)$ and $\tau : \mathcal{A} \to \mathcal{A}$ by

$$\Phi(a,b) = a \text{ and } \tau(a,b) = (b,a).$$

It is clear that $\mathcal{A}$ is a finite set, $\Phi$ and $\tau$ are well-defined, and that $\tau^2 = \mathrm{id}_{\mathcal{A}}$.

Define $\theta : X \to \mathcal{A}^{\mathbb{Z}}$ by

$$\theta(x)_i = \big(x_i\,,\ \varphi(x)_{-i}\big).$$

Since $\varphi$ is (uniformly) continuous, there is a non-negative integer $N$ such that

$$x, x' \in X \text{ and } x_{[-N,\,N]} = x'_{[-N,\,N]} \Rightarrow \varphi(x)_0 = \varphi(x')_0.$$

So, $\theta$ is a sliding block code with memory $N$ and anticipation $N$, and $\theta(X)$ is a shift space over $\mathcal{A}$.

We set $Y = \theta(X)$ and define $\psi : Y \to Y$ by

$$\psi(y)_i = \tau(y_{-i}).$$

It is then straightforward to check that $\psi$ is well-defined, is a flip for $(Y,\, \sigma_Y)$ and that $\Phi_\infty$ is a conjugacy from $(Y,\, \sigma_Y,\, \psi)$ to $(X,\, \sigma_X,\, \varphi)$. □

Now, we prove Proposition 2.1. Let $X$ be a sofic shift and $\varphi$ a flip for $(X,\, \sigma_X)$. With the aid of Lemma 3.1, we may assume that $\varphi$ is a one-block flip. We put $\mathcal{B}_1(X) = \mathcal{B}$ and denote the symbol map of $\varphi$ by $\tau$. Thus $\tau : \mathcal{B} \to \mathcal{B}$, $\tau^2 = \mathrm{id}_{\mathcal{B}}$ and we have

$$\varphi(x)_i = \tau(x_{-i}) \qquad (x \in X,\ i \in \mathbb{Z}).$$

We prove Proposition 2.1 by showing that Krieger's joint state chain of the sofic system $(X, \sigma_X)$ has the desired properties. In the case when $X$ is irreducible(topologically transitive) are further show that Krieger's joint finitary state chain also has the desired properties.

We denote the set of all left-infinite sequences, right-infinite sequences and blocks in $X$ by $L$, $R$ and $B$, respectively:

$$L = \bigcup_{j \in \mathbb{Z}} X_{(-\infty,\, j]},\ R = \bigcup_{j \in \mathbb{Z}} X_{[j,\, \infty)} \text{ and } B = \bigcup_{-\infty < j \le k < \infty} X_{[j,\, k]}.$$

The rules "$\sigma(x)_i = x_{i+1}$" and "$\varphi(x)_i = \tau(x_{-i})$" will be applied to those sequences as well. For instance, if $\lambda \in X_{(-\infty,\, j]}$, then $\sigma(\lambda) \in X_{(-\infty,\, j-1]}$, $\varphi(\lambda) \in X_{[-j,\, \infty)}$,

$$\sigma(\lambda)_i = \lambda_{i+1} \qquad (i \le j-1),$$

and

$$\varphi(\lambda)_i = \tau(\lambda_{-i}) \qquad (i \ge -j).$$

For notational simplicity, we write $\varphi(x) = x^*$ whenever $x \in L \cup R \cup B$. Thus we have $(x^*)^* = x$ for all $x$,

$$L = \{\rho^* : \rho \in R\},\ R = \{\lambda^* : \lambda \in L\} \text{ and } B = \{w^* : w \in B\}.$$

We also write $a^* = \tau(a)$ for $a \in \mathcal{B}$.

If $j, k \in \mathbb{Z}$, $\lambda \in X_{(-\infty,\, j]}$ and $\rho \in X_{[k,\infty)}$, we denote the bi-infinite sequence $x \in \mathcal{B}^{\mathbb{Z}}$ such that $x_{(-\infty,\, -1]} = \sigma^{j+1}(\lambda)$ and $x_{[0,\, \infty)} = \sigma^k(\rho)$ by $\lambda.\rho$ :

$$(\lambda.\rho)_i = \begin{cases} \lambda_{i+j+1} & (i \le -1), \\ \rho_{i+k} & (i \ge 0). \end{cases}$$

We will use similar notation for other sequences in $L \cup R \cup B$ as well.

If $\lambda \in L$, its future is the set $\mathbb{F}(\lambda) \subset X_{[0,\, \infty)}$ defined by

$$\mathbb{F}(\lambda) = \{\rho \in X_{[0,\infty)} : \lambda.\rho \in X\}.$$

The pasts are similarly defined:

$$\mathbb{P}(\rho) = \{\lambda \in X_{(-\infty,\, 0]} : \lambda.\rho \in X\} \qquad (\rho \in R).$$

Is is well known that the sets

$$\Xi_+ = \{\mathbb{F}(\lambda) : \lambda \in L\} \text{ and } \Xi_- = \{\mathbb{P}(\rho) : \rho \in R\}$$

are finite [K]. A triple $(F, a, P) \in \Xi_+ \times \mathcal{B} \times \Xi_-$ is called a joint state of $(X, \sigma_X)$ if $a \in P_0(F) \cap P_0(P)$, where

$$P_0(F) = \{\rho_0 : \rho \in F\} \text{ and } P_0(P) = \{\lambda_0 : \lambda \in P\}.$$

We denote the set of all joint states of $(X, \sigma_X)$ by $\Omega$ and define $\mathcal{L} : \Omega \to \mathcal{B}$ by

$$\mathcal{L}(F, a, P) = a.$$

If $F \in \Xi_+$ and $a \in P_0(F)$, then the set

$$F(a) = \{\rho \in X_{[0,\infty)} : \sigma^{-1}(a.\rho) \in F\}$$

is an elements of $\Xi_+$; and $P \in \Xi_-$ and $a \in P_0(P)$, then the set

$$P(a) = \{\lambda \in X_{(-\infty,0]} : \lambda.a \in P\}$$

is an element of $\Xi_-$. We define the zero-one $\Omega \times \Omega$ matrix $A$ by

$$A((F_1, a_1, P_1), (F_2, a_2, P_2)) = 1 \quad \Leftrightarrow \quad F_2 = F_1(a_1) \text{ and } P_1 = P_2(a_2).$$

The topological Markov chain $\mathsf{X}_A$ is called Krieger's joint state chain of $(X, \sigma_X)$. It is well known that $\mathcal{L}_\infty : \mathsf{X}_A \to X$ is a factoring [K] and it is easy to check that $\mathcal{L}_\infty$ has no graph diamonds.

If $F \in \Xi_+$ and $P \in \Xi_-$, then the sets

$$F^* = \{\rho^* : \rho \in F\} \text{ and } P^* = \{\lambda^* : \lambda \in P\}$$

are elements of $\Xi_-$ and $\Xi_+$, respectively. For $(F, a, P) \in \Omega$ we write

$$(F, a, P)^* = (P^*, a^*, F^*).$$

If $\mathbf{a} \in \Omega$, we have $\mathbf{a}^* \in \Omega$ and $(\mathbf{a}^*)^* = \mathbf{a}$. We also have

$$A(\mathbf{a}, \mathbf{b}) = A(\mathbf{b}^*, \mathbf{a}^*) \qquad (\mathbf{a}, \mathbf{b} \in \Omega).$$

Consequently, if we put $\mathcal{A} = \Omega$ and define the zero-one $\mathcal{A} \times \mathcal{A}$ matrix $J$ by

$$J(\mathbf{a}, \mathbf{b}) = 1 \quad \Leftrightarrow \quad \mathbf{a}^* = \mathbf{b},$$

then the system $(\mathcal{A}, \mathcal{L}, A, J)$ satisfies the condition (1) through (4) in Proposition 2.1.

To prove (5), suppose $x \in X$. Since $\mathcal{L}_\infty : \mathsf{X}_A \to X$ is a factoring, there is a $y \in \mathsf{X}_A$ such that $\mathcal{L}_\infty(y) = x$. We may write

$$y = (F_i, x_i, P_i)_{i \in \mathbb{Z}}.$$

We then have

$$\varphi_{J,A}(y)_i = (P^*_{-i}, x^*_{-i}, F^*_{-i}) \qquad (i \in \mathbb{Z})$$

and

$$\sigma_A \, \varphi_{J,A}(y) = (P^*_{-(i+1)}, x^*_{-(i+1)}, F^*_{-(i+1)}) \qquad (i \in \mathbb{Z}).$$

If $\varphi(x) = x$, then $x_i = (x_{-i})^*$ for all $i$ and hence the point

$$y' = (F_i, x_i, F^*_{-i})_{i \in \mathbb{Z}} \in \mathsf{X}_A$$

satisfies $\mathcal{L}_\infty(y') = x$ and $\varphi_{J,A}(y') = y'$; and if $\sigma_X \, \varphi(x) = x$, then $x_i = (x_{-(i+1)})^*$ for all $i$ and hence the point

$$y'' = (F_i, x_i, F^*_{-(i+1)}) \in \mathsf{X}_A$$

satisfies $\mathcal{L}_\infty(y'') = x$ and $\sigma_A\,\varphi_{J,A}(y'') = y''$. This completes the proof of Proposition 2.1.

A block $w \in B$ is said to be finitary(intrinsically synchronizing) if it has following property:
If $\lambda \in L$, $\rho \in R$, $\lambda.w \in L$ and $w.\rho \in R$, then $(\lambda.w).\rho \in X$.
The set of all finitary blocks in $X$ will be denoted by $B^0$. It is obvious that if $w \in B^0$, then the sets

$$\mathbb{F}(w) = \{\rho \in X_{[0,\infty)} : w.\rho \in R\} \text{ and } \mathbb{P}(w) = \{\lambda \in X_{(-\infty,0]} : \lambda.w \in L\}$$

are elements of $\Xi_+$ and $\Xi_-$, respectively. It is also obvious that $w \in B^0$ if and only if $w^* \in B^0$.

We put

$$\Xi_+^0 = \{\mathbb{F}(w) : w \in B^0\},\ \ \Xi_-^0 = \{\mathbb{P}(w) : w \in B^0\}$$

and

$$\Omega^0 = \{(F, a, P) \in \Omega : F \in \Xi_+^0 \text{ and } P \in \Xi_-^0\}.$$

We denote the resrictions of $\mathcal{L}$, $A$ and $J$ to $\Omega^0$ by $\mathcal{L}^0$, $A^0$ and $J^0$, respectively. The topological Markov chain $\mathsf{X}_{A^0}$ is called Krieger's joint finitary state chain of $(X, \sigma_X)$.

If $X$ is irreducible, it is well known that $(\mathcal{L}^0)_\infty : \mathsf{X}_{A^0} \to X$ is a factoring [K]. Now, it is clear that if $X$ is irreducible, then the system $(\Omega^0, \mathcal{L}^0, A^0, J^0)$ also satisfies the conditions in Proposition 2.1.

## 4. An Example and Some Remarks

**Remark.** Let $X$ be a sofic shift and $\varphi$ a one-block flip for $(X, \sigma_X)$. Suppose $(\mathcal{A}, \mathcal{L}, A, J)$ satisfy the conditons (1) through (4) in Proposition 2.1. Then the conditon (5) is equivalent to the followings:
(i) for any $a \in \mathcal{B}_1(X)$ and $\rho \in F(a)$ with $a^* = a$ and $\rho^*.a\rho \in X$ there exist $b \in \mathcal{L}^{-1}(a)$ and $\lambda \in \mathcal{L}^{-1}(\rho)$ such that $b^* = b$ and $\sigma^{-1}(b.\lambda) \in (\mathsf{X}_A)_{[0,\infty]}$, and
(ii) for any $a \in \mathcal{B}_1(X)$ and $\rho \in F(a)$ with $a^*a \in \mathcal{B}_2(X)$ and $\rho^*a^*.a\rho \in X$ there exist $b \in \mathcal{L}^{-1}(a)$ and $\lambda \in \mathcal{L}^{-1}(\rho)$ such that $b^*b \in \mathcal{B}_2(\mathsf{X}_A)$ and $\sigma^{-1}(b.\lambda) \in (\mathsf{X}_A)_{[0,\infty]}$.

The joint finitary state chain of a irreducible sofic shift may not be irreducible. If we take

$$\Omega' = \{(\mathbb{F}(w_1),\ a,\ \mathbb{P}(w_2)) \in \Omega^0 : w_1 a w_2 \in B\},$$

then $(\Omega', \mathcal{L}', A', J')$ is a irreducible component which satisfies the conditions (1) through (4) in Proposition 2.1, where $\mathcal{L}$, $A'$ and $J'$ are restrictions of $\mathcal{L}$, $A$ and $J$ to $\Omega'$. However, it may not satisfy the condition (5).

For example, the even shift $(X, \sigma)$ with the flip map $\varphi$ induced by the symbol map $\tau = \mathrm{id}_{\{0,1\}}$ has a fixed point $0^\infty \in P(2m, 0)$ for any positive integer $m$, but there is not $y \in (\mathsf{X}_{A'}, \sigma_{A'})$ such that ${\sigma_{A'}}^{2m}(y) = \varphi_{J'A'}(y) = y$. In this case, $(\Omega', \mathcal{L}', A', J')$ dose not satisfy the condition (i) of the above Remark.

**Example.** *Let $(X, \sigma)$ be the even shift, and $\varphi : X \to X$ be the flip induced by the symbol map $\tau = \mathrm{id}_{\{0,1\}}$. Then the zeta fuction is given by*

$$\zeta_{\sigma,\varphi}(t) = \sqrt{\frac{1+t^2}{1-t^2-t^4}}\ \exp\left(\frac{2t+2t^2-t^3-t^4-2t^5-t^6}{(1-t^2)(1-t^2-t^4)}\right).$$


$(\mathbb{F}(1),\ 1,\ \mathbb{P}(1))$
$(\mathbb{F}(1),\ 0,\ \mathbb{P}(01))$
$(\mathbb{F}(10),\ 0,\ \mathbb{P}(1))$
$(\mathbb{F}(1),\ 0,\ \mathbb{P}(1))$
$(\mathbb{F}(10),\ 0,\ \mathbb{P}(01))$

Figure. The joint finitary state chain of the even shift

## References

[B] R. Bowen, *On Axiom A Diffeomorphisms*, AMS-CBMS Reg. Conf. 35, Providence, 1978

[K] W. Krieger, *On Sofic Systems I*, Israel Journal of Mathematics, 48(1984), 305-332.

[KLP] Y.-O. Kim, J. Lee and K. K. Park, *A Zeta Function for Flip Systems*, Pacific Journal of Mathematics, 209(2003), 289-301.

[L] D. Lind, *A zeta function for $\mathbb{Z}^d$-actions*, London Math. Soc. Lecture Note Ser., 228 (M. Pollicott and K.Schmidt, ed.), 1996, 433-450.

[LM] D. Lind and B. Marcus, *Symbolic Dynamics and Coding*, Cambridge University Press, 1995.

[M] A. Manning, *Axiom A diffeomorphisms have rational zeta functions*, Bull. London Math. Soc. 3(1971), 215-220